

Bankart/Instability Rehabilitation Protocol

Ethan M. Healy, M.D.

Post-op Rehabilitation Protocol – SLAP Repair

NO abduction/ER coupled motion for the first 6 weeks post-op

NO biceps resistance exercise for 8 weeks post-op

Weeks 0-4:

Maintain elbow wrist range of motion, Decrease swelling, pain

Treatment plan:

- 1) Participate in pendulum exercises
- 2) Swelling and pain control with ice
- 3) Active elbow, wrist and hand exercises

Weeks 4-8:

Increased shoulder flexibility (limits: Forward Flexion - full, ABD -90, ER -30)

Treatment plan:

- 1) Progressive ROM, and passive stretching (avoid full external rotation till 6 weeks after surgery)
- 2) Swelling and pain control with ice, modalities
- 3) Scapulothoracic strengthening, shoulder isometrics, scapular stabilizers
- 4) Active elbow, wrist and hand exercises

Weeks 8-12:

Full shoulder ROM, progressive strengthening

Treatment plan:

- 1) Continue with swelling and pain control
- 2) Strengthening program: UBE, band weights (start at 2oz-2lbs)
- 3) Rotator cuff/ shoulder PRE
- 4) **Ok to run or jog**

Cont. on next page

Weeks 12-20:

Full shoulder ROM, progressive strengthening

Treatment plan:

- 1) Sports specific training
- 2) Light PNF strengthening
- 3) Rotator cuff/ shoulder PRE
- 4) **Ok to golf**

Weeks 20-30:

Goals: Return to full contact sports if

- 1) pain free
- 2) full ROM
- 3) full strength

Treatment plan:

- 1) Sports specific training
- 2) Light PNF strengthening
- 3) Rotator cuff/ shoulder PRE